SANDY LOAM OVER RED CLAY ON ROCK

General Description: Hard sandy loam to loam abruptly overlying a well structured red clay, calcareous with depth, grading to weathering basement rock.

Landform: Slopes of undulating to

rolling rises or low hills.

Substrate: Fine sandstone or siltstone

> (Appila Formation at this site), mantled by fine

carbonate.

Vegetation:



Type Site: Site No.: CL901 1:50,000 mapsheet: 6629-2 (Kapunda)

Hundred: Easting: 303850 Kapunda 6198150 Section: Northing: 141 Sampling date: 06/03/91 Annual rainfall:

505 mm average

Upper slope of undulating rise, 5% slope. Hard setting surface with about 5% surface siltstone.

Soil Description:

Depth (cm) Description

0-31 Brown hard massive fine sandy loam. Abrupt to:

31-47 Dark reddish brown hard medium clay with

strong medium polyhedral structure. Abrupt to:

47-63 Yellow firm massive very highly calcareous loam

(decomposed rock and 20-50% fine carbonate).

Gradual to:

63-100 Weathering siltstone with 20-50% fine carbonate.

Classification: Sodic, Hypercalcic, Red Chromosol; thick,

slightly gravelly, loamy / clayey, moderate







Summary of Properties

Drainage: Well drained. The soil is unlikely to remain wet for more than a day or so following

heavy or prolonged rainfall.

Fertility: Inherent fertility is moderate. Surface clay and organic matter contents are slightly

low for optimal nutrient retention.

pH: Acidic in the surface, strongly alkaline with depth.

Rooting depth: 100 cm in pit, but few roots below 50 cm.

Barriers to root growth:

Physical: There are no significant barriers above basement rock.

Chemical: High pH in the carbonate layers restricts root growth.

Waterholding capacity: Approximately 80 mm in the rootzone.

Seedling emergence: Fair. Hard setting surface tends to seal over, preventing full seedling emergence.

Workability: Fair. The surface soil tends to shatter if worked too dry, and puddle if worked too

wet.

Erosion Potential:

Water: Moderate.

Wind: Low.

Laboratory Data

Depth cm	pH H ₂ O	pH CaC1 ₂	CO ₃ %	EC1:5 dS/m	ECe dS/m	Org.C %	P	Avail. K mg/kg	mg/kg	Boron mg/kg	Trace Elements mg/kg (DTPA)				CEC cmol (+)/kg	Exchangeable Cations cmol(+)/kg				ESP
											Cu	Fe	Mn	Zn	()	Ca	Mg	Na	K	
0-31	6.2	5.4	0	0.10	1	0.98	47	420	-	-	0.7	37	13.4	1.7	1	1	-	-	-	-
31-47	7.6	6.7	1.8	0.12	1	0.46	5	410	-	3	2.4	11	7.0	0.2	ı	ı	-	-	-	-
47-63	9.1	8.0	43.2	0.14	ı	0.31	1	310	-	1	0.6	1.8	0.4	0.1	1	ı	-	-	-	-
63-100	10.1	8.5	35.6	0.34	-	0.21	1	530	-	-	0.4	2.4	0.5	0.1	1	-	-	-	-	-

Further information: DEWNR Soil and Land Program

