THICK LOAMY SAND OVER MOTTLED BROWN CLAY

General Description: Thick loamy sand over poorly structured brown mottled clay with calcareous subsoil

Landform: Ridge – corridor system of

ancient stranded shoreline.

Substrate: Calcarenite and

lagoonal/fluvial clays.

Vegetation:



Type Site: Site No.: SE024 1:50,000 mapsheet: 7023-1 (Struan)

Hundred:RobertsonEasting:481300Section:376Northing:5894950

Sampling date: 11/03/2005 Annual rainfall: 600 mm average

Footslope of Naracoorte Range, 3% slope. Soft surface with no stones.

Soil Description:

Depth (cm) Description 0-12 Dark brown loamy sand, single grain, loose when dry, water repellent. Sharp break to: 12-45 Very dark greyish brown loamy sand, massive, very firm when dry. Diffuse change to: 45-73 Dark brown loamy sand, massive, very firm when dry. Sharp change to: 73-106 Yellowish brown heavy clay, mottled red, with strong 50-100 mm columnar structure, breaking to strong 20-50 mm polyhedral structure, very firm when moderately moist. Sharp change to: 106-111 Dark brown light medium clay, mottled pinkish white, with strong 20-50 mm polyhedral structure. Much carbonate as soft veins and hard fragments. Sharp break to:

Red heavy clay, massive, very firm when

Red neavy etay, massive, very min when

moderately moist.



Classification: Mottled, Hypercalcic, Brown Chromosol; very thick, non-gravelly, sandy / clayey, deep



111-150



Summary of Properties

Drainage: Imperfect. Water can perch on top of the clayey subsoil for up to several weeks

following heavy or prolonged rainfall. Where topsoil is more than 50 cm thick, perched watertables are unlikely to be a serious problem. Where topsoil is thinner than 30 cm, this duration of waterlogging is significant. Note that topsoil thickness varies from 30 cm to 75 cm in less than a metre across the pit face, so considerable

variability in waterlogging across a paddock is likely.

Fertility: Inherent fertility is moderately low as indicated by the exchangeable cation data. Test

results show marginal phosphorus concentrations and low copper.

pH: Slightly to moderately alkaline throughout. Topsoil alkalinity and detectable

carbonate due to lime dust from adjacent laneway.

Rooting depth: Most root development is in the surface horizons and into the upper clay. Few roots

penetrate into and below the carbonate layer at 106 cm.

Barriers to root growth:

Physical: Compacted hard cap on clayey subsoil restricts root development.

Chemical: There are no apparent chemical barriers.

Waterholding capacity: Approximately 100 mm in the potential rootzone. Varies depending on thickness of

topsoil (refer to profile image).

Seedling emergence: Water repellent surface affects germination. There is potential for sand blasting of

emerging plants.

Workability: Sandy surface is easily worked, but cultivation increases erosion hazard.

Erosion Potential:

Water: Low

Wind: Moderate due to sandy surface.

Laboratory Data

Depth cm	pH H ₂ O	pH CaC1 ₂	CO ₃	EC 1:5 dS/m	ECe dS/m	Org.C	P	Avail. K	Cl mg/kg		Boron mg/kg	Trace Elements mg/kg (EDTA)				Sum cations	Exchangeable Cations cmol(+)/kg				Est. ESP
							mg/kg	mg/kg				Cu	Fe	Mn	Zn	cmol (+)/kg	Ca	Mg	Na	K	
0-12	7.7	7.2	1.0	0.3	3.7	1.1	31	495	124	18.8	0.6	0.57	48	36.4	3.45	10.0	8.09	0.52	0.08	1.33	0.8
12-45	8.0	7.2	0.3	0.1	0.9	0.3	32	126	23	3.5	0.3	0.66	85	31.2	1.05	5.4	4.78	0.20	0.02	0.35	0.4
45-73	8.1	7.0	0.4	0.1	0.3	0.1	24	75	2	2.0	0.3	0.50	90	45.1	0.24	2.9	2.50	0.18	0.01	0.16	na
73-106	7.6	6.6	0.4	0.1	0.2	0.3	3	399	6	2.5	1.0	0.23	62	41.2	0.27	33.3	21.8	9.63	0.48	1.37	1.4
106-111	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
111-150	8.3	7.5	0.7	0.1	0.3	0.3	4	274	5	2.6	2.1	0.24	37	45.4	0.78	26.4	17.4	7.89	0.41	0.75	1.6

Note: Sum of cations, in a neutral to alkaline soil, approximates the CEC (cation exchange capacity), a measure of the soil's capacity to store and release major nutrient elements.

ESP (exchangeable sodium percentage) is derived by dividing the exchangeable sodium value by the CEC, in this case estimated by the sum of cations.

Further information: DEWNR Soil and Land Program



